

APPM 1360 Final Exam Formula Sheet

1. A short table of integrals. In the following, $a \neq 0$.

$$(a) \int \frac{du}{\sqrt{a^2 - u^2}} = \sin^{-1}(u/a) + C \text{ for } u^2 < a^2$$

$$(b) \int \frac{du}{a^2 + u^2} = (1/a) \tan^{-1}(u/a) + C$$

$$(c) \int \frac{du}{u\sqrt{u^2 - a^2}} = (1/a) \sec^{-1} |u/a| + C \text{ for } u^2 > a^2 + C$$

$$(d) \int \frac{du}{\sqrt{a^2 + u^2}} = \sinh^{-1}(u/a) + C \text{ for } a > 0$$

$$(e) \int \frac{du}{\sqrt{u^2 - a^2}} = \cosh^{-1}(u/a) + C \text{ for } u > a > 0$$

$$(f) \int \frac{du}{a^2 - u^2} = \begin{cases} (1/a) \tanh^{-1}(u/a) + C & \text{if } u^2 < a^2 \\ (1/a) \coth^{-1}(u/a) + C & \text{if } u^2 > a^2 \end{cases}$$

$$(g) \int \frac{du}{u\sqrt{a^2 - u^2}} = -(1/a) \operatorname{sech}^{-1}(u/a) + C \text{ for } 0 < u < a$$

$$(h) \int \frac{du}{u\sqrt{a^2 + u^2}} = -(1/a) \operatorname{csch}^{-1} |u/a| + C \text{ for } u \neq 0$$

2. Some trig identities.

$$(a) \sin^2 x + \cos^2 x = 1$$

$$(d) \sin^2 x = (1 - \cos(2x))/2$$

$$(b) \cos^2 x = (1 + \cos(2x))/2$$

$$(e) \cosh^2 x - \sinh^2 x = 1$$

$$(c) \sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$(f) \cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

3. Some useful trig substitutions.

$$x = a \tan \theta \text{ replaces } a^2 + x^2 \text{ by } a^2 \sec^2 \theta$$

$$x = a \sin \theta \text{ replaces } a^2 - x^2 \text{ by } a^2 \cos^2 \theta$$

$$x = a \sec \theta \text{ replaces } x^2 - a^2 \text{ by } a^2 \tan^2 \theta$$

4. Some useful limits.

$$(a) \lim_{n \rightarrow \infty} \frac{\ln n}{n} = 0$$

$$(b) \lim_{z \rightarrow 0} \frac{\sin z}{z} = 1$$

$$(c) \lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$$

$$(d) \lim_{n \rightarrow \infty} \sqrt[n]{\ln(n)} = 1$$

$$(e) \lim_{n \rightarrow \infty} \sqrt[n]{x} = 1 \text{ for } x > 0$$

$$(f) \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x \text{ for any } x$$

$$(g) \lim_{n \rightarrow \infty} x^n = 0 \text{ for } |x| < 1$$

$$(h) \lim_{n \rightarrow \infty} \frac{x^n}{n!} = 0 \text{ for any } x$$

$$(i) \lim_{n \rightarrow \infty} \frac{n!}{n^n} = 0$$

Note: in (e) - (h), x remains **fixed** as $n \rightarrow \infty$.

5. Frequently used Maclaurin series

$$(a) \frac{1}{1-x} = \sum_{n=0}^{\infty} x^n \text{ for } |x| < 1$$

$$(e) e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \text{ for } |x| < \infty$$

$$(b) \sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} \text{ for } |x| < \infty$$

$$(f) \cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} \text{ for } |x| < \infty$$

$$(c) \ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n}, \text{ for } -1 < x \leq 1$$

$$(g) \tan^{-1} x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)}, \text{ for } |x| \leq 1$$

$$(d) (1+x)^m = 1 + \sum_{k=1}^{\infty} \binom{m}{k} x^k \text{ for } |x| < 1$$