


1. (a) (3 pts) Graph the region bounded by $y = x - 2$ and $x = y^2$.
 (b) (12 pts) Find the volume of the solid created by revolving the region bounded by $y = x - 2$ and $x = y^2$ about the line $y = 3$.

Solution:

- (a) The graph




- (b) Here $\Delta V = 2\pi r h$ with $r = 3 - y$ and $h = (y + 2) - y^2$ and so

$$V = \int_{-1}^2 2\pi (3 - y)(y + 2 - y^2) dy = 2\pi \int_{-1}^2 (y^3 - 4y^2 + y + 6) dy = 2\pi \left[\frac{y^4}{4} - \frac{4y^3}{3} + \frac{y^2}{2} + 6y \right]_{-1}^2 = \frac{45\pi}{2}$$

2. (a) (2 pts) Graph the region bounded by the curve $x^3 = 4/y$, and the lines $x = 1$ and $y = 1/2$.
 (b) (8 pts) Set up, but **do not solve**, an integral (or integrals) to find the volume of the solid generated by revolving the region bounded by the curve $x^3 = 4/y$, and the lines $x = 1$ and $y = 1/2$ about the line $x = -e$, using the *disk/washer method*

Solution:

- (a) The graph



- (b) Here $\Delta V = \pi[R^2 - r^2]$ with $R = \sqrt[3]{4/y} + e$ and $r = 1 + e$ and so

$$V = \int_{1/2}^4 \pi \left[\left(\sqrt[3]{4/y} + e \right)^2 - (1 + e)^2 \right] dy$$

-
3. (10 pts) Solve the following differential equations: (a) $2\sqrt{xy}\frac{dy}{dx} = 1$ (b) $\frac{dx}{dt} = e^{t-x}$

Solution:

(a) Separating yields $y^{1/2}dy = \frac{x^{-1/2}}{2}dx$ and so integrating and rearranging terms yields

$$y^{3/2} = \frac{3}{2}x^{1/2} + C, \text{ and so } y = \left(\frac{3}{2}x^{1/2} + C\right)^{2/3}$$

(b) Separating yields $e^x dx = e^t dy$ and so integrating and rearranging terms yields

$$e^x = e^t + C, \text{ and so } x = \ln(e^t + C)$$

-
4. (20 pts) Consider a thin metal plate with density $\rho(x) = 2$ covering the region bounded by the curve $y = \arctan(x)$, $0 \leq x \leq 1$ and the x -axis.

(a) Find the mass of the thin metal plate.

(b) Set-up, **but do not solve**, an integral to find the moment about the x -axis, M_x , of the thin metal plate.

Solution:

(a) Note that here $M = \int \rho(x)dA = \int_0^1 2 \arctan(x)dx$, where this integral is done using integration by parts with $u = \arctan(x)$ and $dv = dx$, then $du = dx/1+x^2$ and $v = x$, so

$$M = \int_0^1 \rho(x)dA = \int_0^1 2 \arctan(x)dx \underset{\text{using I.B.P}}{=} 2x \arctan(x) \Big|_0^1 - \int_0^1 \frac{2x}{1+x^2}dx$$

Now, we use a u -substitution to solve the last integral with $u = 1+x^2$, then $du = 2xdx$ and

$$\int \frac{2x}{1+x^2}dx \underset{u=1+x^2}{=} \int \frac{du}{u} = \ln|u| = \ln(1+x^2)$$

so,

$$M = \int_0^1 2 \arctan(x)dx = 2x \arctan(x) \Big|_0^1 - \ln(1+x^2) \Big|_0^1 = 2 \arctan(1) - \ln(2) = \frac{2\pi}{4} - \ln(2)$$

(b) Here, if we partition the x -axis, then the center of each strip is $(\tilde{x}, \tilde{y}) = (x, \arctan(x)/2)$, thus the moment about the x -axis is given by

$$M_x = \int \tilde{y}dm = \int \tilde{y} \cdot \rho(x)dA = \int_0^1 \frac{\arctan(x)}{2} \cdot 2 \arctan(x)dx = \int_0^1 \arctan^2(x)dx$$

5. (10 pts) Find the length of the curve $y = \cosh(x)$ from $x = 0$ to $x = \ln(2)$.

Solution:

Here $ds = \sqrt{1 + y'(x)^2} dx = \sqrt{1 + \sinh^2(x)} dx$ and so

$$L = \int ds = \int_0^{\ln(2)} \sqrt{1 + \sinh^2(x)} dx \quad \underbrace{=}_{\cosh^2(x)=1+\sinh^2(x)} \int_0^{\ln(2)} \sqrt{\cosh^2} dx = \sinh(x) \Big|_0^{\ln(2)} = \sinh(\ln(2)) = \frac{3}{4}$$

6. (35 pts) Determine if the following converge or diverge. Provide proof of your answers - you must name any test that you use. If any of the series converge, find the sum if possible. (5 pts each)

(a) $\sum_{n=1}^{\infty} \frac{9}{(3n-1)(3n+2)}$ (b) $\left\{ \frac{n + \ln(n)}{n} \right\}_{n=1}^{\infty}$ (c) $\left\{ \frac{n \sin(n)}{3 + n^2} \right\}$ (d) $\sum_{m=2}^{\infty} \frac{e^{2m}}{\pi^m}$

(e) $\{f_m\}_{m=106}^{\infty}$ where $f_m = 1 + (.90)^m$ (f) $\sum_{b=3}^{\infty} \ln(3b) - \ln(2b)$ (g) $\sum_{n=2}^{\infty} \frac{5(-1)^n}{3^{2n}}$

Solution: (a) Using partial fractions we have

$$\frac{9}{(3n-1)(3n+2)} = \frac{A}{3n-1} + \frac{B}{3n+2}, \text{ so } 3A + 3B = 0, \text{ and } 2A - B = 9 \text{ and so } A = 3, B = -3$$

and so,

$$\sum_{n=1}^{\infty} \frac{9}{(3n-1)(3n+2)} = \sum_{n=1}^{\infty} \frac{3}{3n-1} - \frac{3}{3n+2} = \left(\frac{3}{2} - \frac{3}{5}\right) + \left(\frac{3}{5} - \frac{3}{8}\right) + \left(\frac{3}{8} - \frac{3}{11}\right) + \dots = \frac{3}{2} - \lim_{n \rightarrow \infty} \frac{3}{n+2} = \frac{3}{2}$$

thus this is a convergent series which converges to $3/2$.

(b) Taking the limit, we have,

$$\lim_{n \rightarrow \infty} \frac{n + \ln(n)}{n} \stackrel{L'H}{=} \lim_{n \rightarrow \infty} 1 + \frac{1}{n} = 1$$

so the sequence converges to 1.

(c) Note that $-1 \leq \sin(n) \leq 1$, and

$$\frac{-n}{3 + n^2} \leq \frac{n \sin(n)}{3 + n^2} \leq \frac{n}{3 + n^2}$$

and $\lim_{n \rightarrow \infty} \frac{-n}{3 + n^2} = \lim_{n \rightarrow \infty} \frac{n}{3 + n^2} = 0$ (by L'Hopitals rule or dominance of powers), and so by the Squeeze Theorem we have that the sequence converges to 0.

(d) Note that $\frac{e^{2m}}{\pi^m} = \left(\frac{e^2}{\pi}\right)^m$ here we have a Geometric Series with $r = \frac{e^2}{\pi}$ and note that $|r| > 1$ and so this is a divergent series.

(e) Taking the limit we have $\lim_{m \rightarrow \infty} f_m = \lim_{m \rightarrow \infty} 1 + (0.9)^m \stackrel{=}{\underbrace{\text{since } 0.9 < 1}} 1$, so the sequence converges to 1.

(f) Note that $\lim_{b \rightarrow \infty} a_b = \lim_{b \rightarrow \infty} \ln(3b) - \ln(2b) = \lim_{b \rightarrow \infty} \ln\left(\frac{3b}{2b}\right) = \ln\left(\frac{3}{2}\right) \neq 0$, and so the series diverges by the Divergence Test

(g) Here we have a geometric with,

$$\sum_{n=2}^{\infty} \frac{5(-1)^n}{3^{2n}} = \sum_{n=2}^{\infty} 5 \left(-\frac{1}{3^2}\right)^n = \frac{5}{81} + \frac{5}{81} \left(-\frac{1}{9}\right) + \frac{5}{81} \left(-\frac{1}{9}\right)^2 + \dots$$

so $a = 5/81$ and $r = -1/9$ and $|r| < 1$ and so

$$\sum_{n=2}^{\infty} \frac{5(-1)^n}{3^{2n}} = \frac{5/81}{1 - (-1/9)} = \frac{5}{90} = \frac{1}{18}$$

thus this is a convergent series which converges to $1/18$.
