

Homework set 13 — APPM5440, Fall 2009

Note: This homework covers all of Chapter 6.

From the textbook: 6.1, 6.2, 6.3, 6.4, 6.6, 6.12.

Problem 1: Let H be a Hilbert space, and let $(e_j)_{j=1}^n$ be an orthonormal set in H . Let x be an arbitrary vector in H . Set $M = \text{span}(e_1, \dots, e_n)$, set

$$y = \sum_{j=1}^n (e_j, x) e_j,$$

and set $z = x - y$. Prove that $z \in M^\perp$ (and consequently, that $y \perp z$). Prove that

$$\|x - y\| = \inf_{y' \in M} \|x - y'\|.$$

Prove that y is the *unique* minimizer (in other words, if $y' \in M \setminus \{y\}$, then $\|x - y'\| > \|x - y\|$). Prove these claims directly, without using the theorem about existence of a unique minimizer between a closed convex set and a point.

Problem 2: Set $I = [-1, 1]$ and consider the Hilbert space $H = L^2(I)$. Let M denote the subspace of H consisting of all even functions (in other words, functions such that $f(x) = f(-x)$ for all x). Given an $f \in H$, prove that

$$\inf_{g \in M} \|f - g\| = \left(\int_{-1}^1 \left(\frac{f(x) - f(-x)}{2} \right)^2 dx \right)^{1/2}.$$

(Don't worry about issues relating to Lebesgue integration.)

The last problem was the “hard” problem on the final last year.

Problem 3: Let X be a separable infinite-dimensional Hilbert space. Prove that there exists a family of closed linear subspaces $\{\Omega_t : t \in [0, 1]\}$ such that Ω_s is a strict subset of Ω_t whenever $s < t$.