

Problem 11.22: Set $T = \text{sign}(t)$. We seek to prove that $\check{T} = \alpha \text{PV}(1/x)$ for some α .

For $N = 1, 2, 3, \dots$, set $T_N = \chi_{[-N, N]} T$. Then $T_N \rightarrow T$ in \mathcal{S}^* since for any $\varphi \in \mathcal{S}$, we have

$$\langle T_n, \varphi \rangle = \int_{-N}^N \text{sign}(x) \varphi(x) dx \rightarrow \int_{-\infty}^{\infty} \text{sign}(x) \varphi(x) dx = \langle T, \varphi \rangle.$$

Since the Fourier transform is a continuous operator on \mathcal{S}^* , we know that \check{T} is the limit of the sequence $(\check{T}_N)_{N=1}^{\infty}$.

Since $T_N \in L^1$, we can compute \check{T}_N by directly evaluating the integral. We find that

$$(1) \quad \check{T}_N(x) = \beta \frac{1 - \cos(Nx)}{x}$$

for some constant β . If $\varphi \in \mathcal{S}$, then

$$\begin{aligned} \left\langle \frac{1 - \cos(Nx)}{x}, \varphi \right\rangle &= \int_{\mathbb{R}} \frac{1 - \cos(Nx)}{x} \varphi(x) dx = \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \frac{1 - \cos(Nx)}{x} \varphi(x) dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \frac{1}{x} \varphi(x) dx - \lim_{\varepsilon \rightarrow 0} \int_{|x| \geq \varepsilon} \cos(Nx) \frac{1}{x} \varphi(x) dx \\ &= \langle \text{PV}(1/x), \varphi \rangle - \langle \cos(Nx) \text{PV}(1/x), \varphi \rangle. \end{aligned}$$

It follows that formula (1) can be written $\check{T}_N(x) = \beta \text{PV}(1/x) - \beta \cos(Nx) \text{PV}(1/x)$.

It remains to prove that $\cos(Nx) \text{PV}(1/x) \rightarrow 0$ in \mathcal{S}' . We find that

$$\begin{aligned} \langle \cos(Nx) \text{PV}(1/x), \varphi \rangle &= \langle \text{PV}(1/x), \cos(Nx) \varphi \rangle \\ &= \int_0^{\infty} \cos(Nx) \frac{1}{x} \varphi(x) dx + \int_{-\infty}^0 \cos(Nx) \frac{1}{x} \varphi(x) dx \\ &= \int_0^{\infty} \cos(Nx) \frac{\varphi(x) - \varphi(-x)}{x} dx. \end{aligned}$$

Now set $\psi(x) = \frac{\varphi(x) - \varphi(-x)}{x}$. Then ψ is a continuously differentiable, quickly decaying function on $[0, \infty)$, so we can perform a partial integration to obtain

$$\begin{aligned} \left| \int_0^{\infty} \cos(Nx) \frac{\varphi(x) - \varphi(-x)}{x} dx \right| &= \left| \left[\frac{\sin(Nx)}{N} \psi(x) \right]_0^{\infty} - \int_0^{\infty} \frac{\sin(Nx)}{N} \psi'(x) dx \right| \\ &\leq \frac{1}{N} \int_0^{\infty} |\psi'(x)| dx. \end{aligned}$$

If we can prove that $\int_0^{\infty} |\psi'(x)| dx < \infty$, we will be done. First note that for $x \in [0, 1]$, $\psi(x) = 2\varphi'(0) + O(x^2)$, so for $x \in [0, 1]$, we have $|\psi'(x)| \leq C_1$ for some finite C_1 . For $x \in [1, \infty)$, we have

$$|\psi'(x)| = \left| \frac{\varphi'(x) + \varphi'(-x)}{x} - \frac{\varphi(x) - \varphi(-x)}{x^2} \right| \leq 2 \frac{\|\varphi\|_{1,1}}{x^2} + 2 \frac{\|\varphi\|_{0,0}}{x^2} = \frac{C_2}{x^2}.$$

and so

$$\int_0^{\infty} |\psi'(x)| dx \leq \int_0^1 C_1 dx + \int_1^{\infty} \frac{C_2}{x^2} dx < \infty.$$