

### Homework set 3 — APPM5450, Spring 2017

From the textbook: 8.6. Optional: 8.5.

**Problem 1:** Let  $H$  be a Hilbert space, and let  $(\varphi_n)_{n=1}^{\infty}$  denote an orthonormal basis for  $H$ . Given a bounded sequence of complex number  $(\lambda_n)_{n=1}^{\infty}$ , define the operator  $A$  by setting

$$A u = \sum_{n=1}^{\infty} \lambda_n \varphi_n \langle \varphi_n, u \rangle.$$

(a) Prove that  $\|A\| = \sup_n |\lambda_n|$ .

(b) Prove that  $A^*u = \sum_{n=1}^{\infty} \bar{\lambda}_n \varphi_n \langle \varphi_n, u \rangle$ . Conclude that  $A$  is self-adjoint iff all  $\lambda_n$ 's are real. When is  $A$  skew-symmetric? When is  $A$  non-negative / positive / coercive?

**Problem 2:** Consider the Hilbert space  $H = L^2([-\pi, \pi])$ , and the operator  $A \in \mathcal{B}(H)$  defined by  $[Au](x) = |x| u(x)$ . Prove that  $A$  is self-adjoint and positive, but not coercive. Prove that

$$\langle u, v \rangle_A = \langle Au, v \rangle$$

is an inner product on  $H$ , but that the topology generated by (the norm generated by) this inner product is *not* equivalent to the topology generated by the  $L^2$ -norm.

**Problem 3:** Set  $H = \ell^2(\mathbb{Z})$  and let  $R$  denote the right-shift operator (so that if  $y = Rx$ , then  $y_n = x_{n-1}$ ). Construct  $R^*$ . Prove that  $RR^* = R^*R = I$ , which is to say that  $R$  is “unitary.” (Is either the right or the left-shift operator on  $\ell^2(\mathbb{N})$  unitary?)

**Problem 4:** Consider the Hilbert space  $L^2(\mathbb{T})$ . Let  $k$  denote a continuous function on  $\mathbb{T}^2$  that takes on complex values. Let  $A$  denote the operator  $[Au](x) = \int_{\mathbb{T}} k(x, y) u(y) dy$ . Prove that  $[A^*u](x) = \int_{\mathbb{T}} \overline{k(y, x)} u(y) dy$ . Conclude that  $A$  is self-adjoint iff  $k(x, y) = \overline{k(y, x)} \forall x, y \in \mathbb{T}$ .