

The removal of critical slowing down

M. A. Clark*

Center for Computational Science, Boston University, 3 Cummington St, MA 02215, USA

E-mail: mikec@bu.edu

J. Brannick^a, R. C. Brower^{bc}, S. F. McCormick^e, T. A. Manteuffel^e, J. C. Osborn^{bd} and C. Rebbi^{bc}

^a *Department of Mathematics, The Pennsylvania State University, 221 McAllister Building, University Park, PA 16802, USA*

^b *Center for Computational Science, Boston University, 3 Cummington St, MA 02215, USA*

^c *Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, MA 02215, USA*

^d *Argonne Leadership Computing Facility, 9700 S. Cass Avenue, Argonne, IL 60439, USA*

^e *Department of Applied Mathematics, Campus Box 526, University of Colorado at Boulder, Boulder, CO 80309, USA*

We present promising initial results of our adaptive multigrid solver developed for application directly to the non-Hermitian Wilson-Dirac system in 4 dimensions, as opposed to the solver developed in [1] for the corresponding normal equations. The key behind the success of this algorithm is the use of an adaptive projection onto coarse grids that preserves the near null space of the system matrix. We demonstrate that the resulting algorithm has weak dependence on the gauge coupling and exhibits extremely mild critical slowing down in the chiral limit.

The XXVI International Symposium on Lattice Field Theory

July 14-19 2008

Williamsburg, Virginia, USA

*Speaker.

1. Introduction

We aim to solve the system of equations

$$Dx = b,$$

where, in general, $D = \mathbb{D} + m$ is the Dirac operator, with \mathbb{D} denoting the tensor product of the discretized covariant derivatives and the Dirac gamma matrices and m the fermion mass. Herein, we restrict our discussion to the Wilson-Dirac discretization of D , obtained by using central covariant differences to discretize D and then adding a properly scaled second-order Wilson term. It is well known that for this choice, D is a non-Hermitian matrix, with complex eigenvalues which, in general, satisfy $\text{Re}(\lambda) > 0$ and also that D satisfies a γ_5 Hermiticity, i.e., $D^\dagger = \gamma_5 D \gamma_5$, and $H = \gamma_5 D$, where H is the Hermitian (indefinite) Dirac operator. As the fermion mass is reduced to zero ($\text{Re}(\lambda_{\min}) \rightarrow 0$), D becomes singular, causing the "critical slowing down" (deteriorating convergence) of the standard Krylov solvers typically used to solve these systems. Improving the convergence of Krylov methods for the Dirac inversion problem with a suitable preconditioning has been a main topic of research in lattice QCD for many years and, until more recently, the cost of inverting D has been prohibitive as this physical light quark mass is approached.

In the last few years, much progress has been made in addressing this issue. Eigenvector deflation [2, 3] is a proven technique for accelerating convergence, provided sufficiently many eigenvectors are used in the deflation; exact deflation approaches are, however, expected to scale as the square of the lattice volume $O(V^2)$ and, thus, become ineffective for large volume. An alternative approach is given by the inexact local mode deflation proposed by Lüscher [4]; here, only approximate eigenvectors are used in the deflation process and, due to the local coherence (see below) exhibited by the low (small magnitude) eigenmodes of the Dirac operator, only a small number of low mode prototypes are needed for an effective deflation. Inexact deflation strategies are expected to scale as $O(V \log V)$ or even $O(V)$.

The approach we are proposing is an alternative to deflation, based on a multigrid (MG) solver for the Dirac operator. In previous work [1], we presented an algorithm for the normal equations obtained from the Wilson-Dirac system in the context of 2 dimensions, with a $U(1)$ gauge field. Here, we extend the approach directly to the Wilson-Dirac system (as opposed to the normal equations) and apply the resulting algorithm to the full 4-dimensional $SU(3)$ problem.

2. Adaptive Multigrid

The "light" modes, low eigenmodes of the system matrix, typically cause the poor convergence suffered by standard iterative solvers (relaxation or Krylov methods); as the operator becomes singular, the error in the iteratively computed solution quickly becomes dominated by these modes. In free field theory, these slow-to-converge modes are locally geometrically smooth and, hence, they can be well represented on a coarse-grid using fewer degrees of freedom. Moreover, these smooth modes on the fine grid now again become rough (high frequency) modes, on the coarse-grid. It is this observation that motivates the classical geometrical MG approach, in which simple local averaging and linear interpolation can be used to form corrections to the fine grids stemming from solutions upon a coarse coarse-grid. We hereafter denote the interpolation operator by P

and restriction operator by R . Given a Hermitian positive definite (HPD) operator A , taking the restriction operator as $R = P^\dagger$ and the coarse-grid operator as $A_c = P^\dagger A P$, gives the optimal, in an energy-norm sense, two-level correction. It is natural to recurse upon this approach by defining the problem on coarser and coarser grids until the degrees of the freedom have been reduced enough to permit an exact solve. When combined with m and n relaxation applications before and after each restriction and prolongation applications, to remove the high frequency error inherent to that level, this is known as a $V(m, n)$ -cycle, and removes critical slowing down for discretized PDE problems and scales as $O(V \log V)$ or $O(V)$ [5].

The error propagation operator for the two-level solver with a single post-relaxation, with error propagator S , is given by

$$E_{TG} = S(I - \pi_A), \quad \pi_A = P(P^\dagger A P)^{-1} P^\dagger A = P A_c^{-1} P^\dagger A.$$

Roughly speaking, the performance of a given MG algorithm is related to $\text{span}(P P^\dagger)$ and how well this approximates the slow-to-converge modes of the chosen relaxation procedure (called a ‘‘smoother’’). More precisely, given P and a convergent smoother, the two-level algorithm can be shown to converge provided: for any fine-level vector $u \in \mathbb{C}^n$, there exists a vector of coefficients $w_c \in \mathbb{C}^{n_c}$ on the coarse-level such that the linear combination $P w_c$, with the columns of P denoting the coarse-space basis functions, satisfies

$$K(P) = \|A\| \frac{\|u - P w_c\|}{\|u\|_A} \leq \infty.$$

The above inequality, known as the weak approximation property in the MG literature, is clearly biased towards the light modes and in particular suggests that P approximate eigenvectors with error proportional to the size of their corresponding eigenvalues.

When solving the Wilson-Dirac system in the interacting theory, the light modes are not geometrically smooth, and so classical MG, that assumes the slow-to-converge error to be locally constant, fails completely. In such settings, where the gauge field is essentially random and this randomness dictates the local nature of the low modes, we must alter our definition of the prolongator P , to span this space. As the columns of P form the coarse space basis, this is achieved by partitioning the light modes into subvectors over the aggregates. The above approximation property then implies that locally the low modes used in defining P must form a basis for the low modes of the system matrix, which for most simple pointwise smoothers are also the low modes not effectively treated by the relaxation method. In the physics community, this notion that a small set of vectors partitioned into local basis functions can approximate the entire lower end of the eigen-spectrum of a matrix is known as local coherence. It is this idea of local coherence that leads to the success of Lüscher’s [4] deflation approach as well as our MG solver.

With a convergent MG algorithm defined, we are now left with the problem of adaptively finding a representation of the light modes to allow us to define P . One viable approach, known in the MG literature as adaptive smooth aggregation (α SA) [6], is given by iteratively computing the low modes and then adjusting P to fit these computed modes. The general algorithm for computing these prototypes for some matrix A proceeds as follows:

1. Apply the MG relaxation to the system $Ax = 0$, where x_0 is taken to be a random vector.

2. Terminate the solver after l iterations. The current iterate $x_l = -e_l = v_1$ will be a representation of the slow to converge components.
3. Decompose this error vector into aggregates (blocks). The components within each of these blocks represent a column of the prolongator.

Numerical experience suggests that for the 4-dimensional Wilson-Dirac system with disordered background gauge field, more than a single near null space vector is needed in defining a P that sufficiently captures the near null space of the operator in question, that is, more than a single vector is needed to ensure local coherence. In the above discussion, the initial prototype of the null space used in defining P is computed using relaxation. More generally, the current MG solver is applied to $Ax = 0$ to generate prototypes of the algebraically smooth error, yielding at each adaptive step the matrix $V^k = [v_1, \dots, v_k]$, with the v_i 's denoting the computed candidate vectors approximating the near kernel of the operator A , where V^k is augmented until convergence is deemed sufficient, say for $k = N_v$ candidate vectors. At each stage in which V^k is augmented, one defines the (tentative) prolongation operator P and the representation of V^k on the coarse lattice V_c^k in the usual way, that is, by enforcing the following relations:

$$PV_c^k = V^k$$

and

$$P^\dagger P = I,$$

that is to say one performs a $\mathcal{Q}\mathcal{R}$ decomposition on the partitioned candidate vectors, where \mathcal{Q} represents the columns of prolongator and \mathcal{R} represents the coefficients in the coarse basis, i.e., V_c^k .

3. Formulating an algorithm

The original adaptive smoothed aggregation approach introduced in [6] is essentially a black-box method, where the blocking strategy is chosen using an algebraic strength-of-connection measure. In lattice QCD, the system is discretized on an uniform hypercubic lattice, with unitary connections link the sites. With this in mind, our blocking strategy is to use a regular geometric structure, e.g., 4^4 geometric blocking.

In solving the Wilson-Dirac system, we consider two approaches:

1. Apply the adaptive MG approach to the normal equations.
2. Formulate a MG algorithm directly for the Wilson-Dirac operator.

The normal equation approach has the obvious advantage that the operator in question is HPD; hence, the standard Galerkin operator definition $(D^\dagger D)_c = P^\dagger (D^\dagger D)P$ is optimal and all MG convergence proofs can be applied directly. However, the condition number of $D^\dagger D = H^2$ is the square of that of H and the coarse operator cannot be written as the product of nearest neighbour couplings, leading to fill-in in the operator.

For non-HPD systems, the usual MG convergence proofs generally do not apply. However, in this case, a significant amount of insight is obtained by considering the spectral decomposition

of $D = |\tilde{\psi}_\lambda\rangle\lambda\langle\psi_\lambda|$, where ψ and $\tilde{\psi}$ are the right and left eigenvectors respectively, both having eigenvalue λ . If we now consider eigenvector deflation, in which case we use a Petrov-Galerkin oblique projection, i.e., treat the left and right spaces separately, to remove a given eigenvalue λ , then we have:

$$\mathcal{P} = \left(1 - D|\psi_\lambda\rangle\frac{1}{\lambda}\langle\tilde{\psi}_\lambda|\right) = (1 - D|\psi_\lambda\rangle\langle\tilde{\psi}_\lambda|D|\psi_\lambda\rangle^{-1}\langle\tilde{\psi}_\lambda|) = (1 - DP(RDP)^{-1}R).$$

We thus see that prolongation should be defined using ‘‘right null space vectors’’ and restriction from ‘‘left null space vectors’’. Naively, this suggests that we define prolongation using smoothed vectors of D and restriction from smoothed vectors of D^\dagger . However, because of the γ_5 symmetry of the Wilson-Dirac operator, we have $\psi_\lambda = \gamma_5\tilde{\psi}_{\lambda^*}$; and hence, a vector rich in low right eigenvectors can be converted to one rich in low left eigenvectors by simple multiplication by γ_5 . Given the current residual r_0 , our coarse-grid correction is thus given by

$$x_c = \alpha P(P^\dagger\gamma_5DP)^{-1}P^\dagger\gamma_5r_0 = P(P^\dagger HP)^{-1}P^\dagger\gamma_5r_0,$$

where the step-size α will be defined below. If we block the spin dimension, however, it is possible for the coarse operator to have exactly zero eigenvalues¹. This is easy to avoid by keeping chirality intact, i.e., $[\gamma_5, P] = 0$. In this way, the γ_5 factors cancel out in the overall coarse-level correction, yielding the former ‘‘naive’’ result $R = P^\dagger$.

The convergence of our coarse-grid correction is not guaranteed: if we perform an eigenvector expansion of the low modes used to define the prolongator

$$v_i = \sum_{\lambda, \lambda^*} \langle\psi_\lambda|v_i\rangle\psi_\lambda + \langle\psi_{\lambda^*}|v_i\rangle\psi_{\lambda^*} = \sum_{\lambda, \lambda^*} c_\lambda\psi_\lambda + c_{\lambda^*}\psi_{\lambda^*},$$

the left space is given by

$$\hat{v}_i = \sum_{\lambda, \lambda^*} c_\lambda\hat{\psi}_{\lambda^*} + c_{\lambda^*}\hat{\psi}_\lambda.$$

Since in general $c_\lambda \neq c_{\lambda^*}$, the left and right spaces are not equivalent, and our coarse-grid correction can be divergent. Convergence can be imposed by choosing the step-size α to always minimize the resulting residual $r_1 = r_0 - Dx_c$, i.e.,

$$\alpha = \frac{\langle Dx_c|r_0\rangle}{\langle Dx_c|Dx_c\rangle}.$$

With the prolongator and coarse-grid operator defined, all that remains is to define a suitable relaxation procedure that effectively damps the eigenvectors of the system matrix with large eigenvalues, in magnitude. Classical MG methods use either Jacobi or Gauss-Seidel smoothing, which are either inefficient or cannot be applied directly to non-HPD operators. Currently, we use an under-relaxed minimal residual smoother (with under-relaxation parameter $\omega = 0.8$). This yields a simple parallel approach that reduces the residual in the $D^\dagger D$ norm, ensuring that error components corresponding to eigenvectors with large eigenvalues are quickly damped.

¹Take for example the free field operator, where the null space vector is given by the constant. Thus, $P^\dagger\gamma_5P = 0$, and our coarse-grid correction is ill-defined.




Figure 1: Comparison of CG, MG-CG ($N_v = 8$) and MG-BiCGstab ($N_v = 3$) algorithms: left panel compares total number of Wilson matrix-vector operations; right panel compares actual flops ($V = 128^2$, $\beta = 6.0$).

4. 2d Results

Before presenting results of our solver to the full $4d$ problem, we compare the normal equations and direct approaches in the $2d$ QED context. Here, a 3-level $V(2,2)$ -cycle approach has been used, with 4^2 geometric blocking and a machine precision solve has been used on the coarsest ($V = 8^2$) lattice. The normal equation approach is used to precondition CG (MG-CG), and the direct approach to precondition BiCGstab (MG-BiCGstab). In the left plot of figure 1, which compares the total number of Dirac operator applications, we see that both approaches effectively remove the critical slowing down, with both methods leading to around an order of magnitude reduction of operator applications. Looking at the actual number of floating point operations (right panel), it is seen that MG-BiCGstab is around an order of magnitude cheaper than MG-CG as the chiral limit is taken. This large disparity in cost between the two approaches is due to the increased operator complexity for the coarse normal equation operator ($N_v = 8$) compared to the direct approach ($N_v = 3$). It should be noted that we have tested the validity of our algorithm in the range $\beta = 0.1 \dots 100$ and found very little dependence with the resulting MG performance.

5. 4d Results

Given the $2d$ results, we applied only the direct approach to the $4d$ operator; here we found that $N_v = 20$ vectors were required to sufficiently capture the null space of the Dirac operator (this matches Lüscher's result in [4]). The outer Krylov solver was switched to GCR(50) since our MG operator is a non-stationary solver (this leads to breakdown in the recursion relations of non-restarted solvers). In figure 2 we demonstrate the benefit of our MG-GCR algorithm over CG and BiCGstab, where our MG algorithm is a modified 3-level $V(2,2)$ -cycle: between each restriction and prolongation application on the fine grid there are 4 applications of the next coarsest level V -cycle, this ensures much more accurate solutions are obtained to the first coarse-grid. Again 4^d geometric blocking is used and in moving from the original fine grid to the first coarse-grid




Figure 2: Comparison of CG, BiCGstab and MG-GCR algorithms, left panel compares total number of Wilson matrix-vector operations, right panel compares time to solution ($V = 16^3.32$, $\beta = 6.0$, $m_{crit} = -0.8049$, $N_v = 20$).

we also include colour and spin components with like chirality in the block (but not the full spin dimension). On the left panel we plot the total number of Dirac operator applications until convergence, where we see that there is an order of magnitude improvement over BiCGstab. On the right panel, comparing time to convergence, we see that the true benefit is closer to a factor of 4-5 over BiCGstab as the chiral limit is taken.

6. Conclusion

In this work, we introduced a new multigrid algorithm and showed that it removes the critical slowing down as the quark mass is taken to zero in the Wilson-Dirac operator.

Future work in this area will focus on further refining our algorithm to reduce the setup cost of the algorithm, i.e., attempt to reduce from $N_v = 20$ and to apply these techniques to chiral fermions.

This research was supported under: DOE grants DE-FG02-91ER40676, DE-FC02-06ER41440, DE-FG02-03ER25574 and DE-FC02-06ER25784; Lawrence Livermore National Laboratory contracts B568677, B574163 and B568399; NSF grants PHY-0427646, DMS-0749317, 0810982 and 0749202.

References

- [1] J. Brannick, R. C. Brower, M. A. Clark, J. C. Osborn and C. Rebbi, Phys. Rev. Lett. **100** (2008) 041601 [arXiv:0707.4018 [hep-lat]].
- [2] A. Stathopoulos and K. Orginos, arXiv:0707.0131 [hep-lat].
- [3] R. B. Morgan and W. Wilcox, arXiv:0707.0505 [math-ph].
- [4] M. Luscher, JHEP **0707**, 081 (2007) [arXiv:0706.2298 [hep-lat]].
- [5] A. Brandt. *Math. Comp.*, 31:333–390, 1977.
- [6] M. Brezina, R. Falgout, S. MacLachlan, T. Manteuffek, S. McCormick, and J. Ruge. *Siam J. Sci. Comput.*, 25:1896–1920, 2004.