

## Homework set 8 — APPM5440 — Fall 2012

**From the textbook:** 3.6, 3.7.

On the next page, you'll find the 2005 midterm. Problem 4 on that midterm is part of this week's homework. (Note that the questions on topological spaces are outside the syllabus for the midterm this year.)

**Problem 1:** Let  $X$  be a set with infinitely many members. We define a collection  $\mathcal{T}$  of subsets of  $X$  by saying that a set  $\Omega \in \mathcal{T}$  if either  $\Omega^c = X \setminus \Omega$  is finite, or if  $\Omega$  is the empty set. Verify that  $\mathcal{T}$  is a topology on  $X$ . This topology is called the “co-finite” topology on  $X$ . Describe the closed sets.

**Problem 2:** Let  $X$  denote a finite set, and let  $\mathcal{T}$  be a metrizable topology on  $X$ . Prove that  $\mathcal{T}$  is the discrete topology on  $X$ .

**Problem 3:** Consider the set  $X = \{a, b, c\}$ , and the collection of subsets  $\mathcal{T} = \{\emptyset, \{a\}, \{a, b\}, \{a, c\}, \{a, b, c\}\}$ . Is  $\mathcal{T}$  a topology? Is  $\mathcal{T}$  a metrizable topology?

**Applied Analysis (APPM 5440): Midterm 2**  
5.00pm – 6.30pm, Oct 26, 2005. Closed books.

**Problem 1:** State the Arzelà-Ascoli theorem.

**Problem 2:** We describe two mathematical objects below. For each description, either provide an example of such an object, or explain why it does not exist. (Brief answers, please!)

- (1) A collection of functions in  $C(\mathbb{R})$  that is equicontinuous but not uniformly equicontinuous.
- (2) A collection of functions in  $C([0, 1])$  that is equicontinuous but not uniformly equicontinuous.

**Problem 3:** Let  $X = \{a, b, c, d\}$  be a set, and let

$$\mathcal{S} = \{\emptyset, X, \{b, d\}, \{a, c\}, \{d\}, \{a, b, c\}\}$$

be a collection of subsets of  $X$ .

- (1) Prove that  $\mathcal{S}$  is not a topology on  $X$ .
- (2) Let  $\mathcal{T}$  denote the smallest topology on  $X$  that contains  $\mathcal{S}$  (in other words,  $\mathcal{T}$  is the topology generated by the sub-base  $\mathcal{S}$ ). List the sets that are contained in  $\mathcal{T}$  but not in  $\mathcal{S}$ .

**Problem 4:** Consider the integral equation

$$(*) \quad u(x) = \pi^2 \sin(x) + \frac{3}{2} \int_0^{\cos(x)} |x - y| u(y) dy.$$

Prove that (\*) has a unique solution in  $C([0, 1])$ .

**Problem 5:** Let  $X$  be a topological space, let  $Y$  be a Hausdorff space, and let  $f$  and  $g$  be continuous maps from  $X$  to  $Y$ . Is it necessarily the case that the set  $\Omega = \{x \in X : f(x) = g(x)\}$  is closed? Justify your answer by either giving a proof or a counter example.